Genomic identification, annotation, and comparative analysis of Vacuolar-type ATP synthase subunits in *Diaphorina citri*

Rebecca Grace^{1,2}, Crissy Massimino¹, Teresa D. Shippy³, Will Tank³, Prashant S. Hosmani⁴, Mirella Flores-Gonzalez⁴, Lukas A. Mueller⁴, Wayne B. Hunter⁵, Joshua B. Benoit⁶, Susan J. Brown³, Tom D'Elia¹ and Surya Saha^{4,7,*}

¹ Indian River State College, Fort Pierce, FL 34981, USA

² Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853, USA

³ Division of Biology, Kansas State University, Manhattan, KS 66506, USA

⁴ Boyce Thompson Institute, Ithaca, NY 14853, USA

⁵ USDA-ARS, US Horticultural Research Laboratory, Fort Pierce, FL 34945, USA

⁶ Department of Biological Sciences, University of Cincinnati, Cincinnati, OH 45221, USA

⁷ Animal and Comparative Biomedical Sciences, University of Arizona, Tucson, AZ 85721, USA

ABSTRACT

The hemipteran insect *Diaphorina citri*, or Asian citrus psyllid, is a vector for *Candidatus Liberibacter asiaticus* (CLas), the bacterium causing citrus greening disease, or Huanglongbing (HLB). Millions of citrus trees have been destroyed, and every grove in Florida, USA, has been directly affected by this disease. In eukaryotes, vacuolar-type ATP synthase (V-ATPase) is an abundant heterodimeric enzyme that serves the cell with essential compartment acidification through the active processes that transport protons across the membrane. Fifteen putative V-ATPase genes in the *D. citri* genome were manually curated. Comparative genomic analysis revealed that *D. citri* V-ATPase subunits share domains and motifs with other insects, including the V-ATPase-A superfamily domain. Phylogenetic analysis separates *D. citri* V-ATPase subunits into expected clades with orthologous sequences. Annotation of the *D. citri* genome is a critical step towards developing directed pest management strategies to reduce the spread of HLB throughout the citrus industry.

Subjects Genetics and Genomics, Animal Genetics, Bioinformatics

INTRODUCTION

Vacuolar (H⁺)-ATP synthase (V-ATPase) is a highly conserved eukaryotic enzyme [1]. Originally identified in the vacuole membrane, V-ATPase has a critical function in the plasma membrane and endomembrane system of almost every cell [2, 3]. V-ATPase regulates the acidity of organelles, such as vacuoles, the Golgi apparatus, and coated vesicles, by translocating protons across their membranes and powering secondary transport processes. Structurally, V-ATPase has a noncatalytic transmembrane domain, the V₀ rotor, and a catalytic cytoplasmic domain, the V₁ stator. V-ATPase hydrolyzes adenosine triphosphate (ATP) into adenosine diphosphate (ADP), thus acting in the opposite manner to

Submitted: 21 October 2021
Accepted: 04 February 2022
Published: 09 February 2022

* Corresponding author. E-mail: suryasaha@cornell.edu

Published by GigaScience Press.

Preprint submitted at <https://doi.org/10.1101/2021.10.18.464890>

Included in the series: **Asian citrus psyllid community annotation** (https://doi.org/10.46471/GIGABYTE_SERIES_0001)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Gigabyte, 2022, 1–18

the related F-ATPase [1]. In insects, 13 protein subunits are typically required to build a single V-ATPase [4]. The V_0 domain comprises subunits a–e, and V_1 comprises subunits A–H [5]. The critical accessory subunit S1 (Ac45) also helps assemble the enzyme [6].

CONTEXT

In insects, high levels of V-ATPase are found in epithelial cells. They are especially important in the digestive tract, helping to regulate nutrient uptake and solute transport [7]. Studies in several phyla, including insects, have demonstrated the lethality of silencing individual *V-ATPase* genes, making *V-ATPase* an attractive target for RNA interference (RNAi)-based pest control [1, 4, 7]. We have characterized the genes encoding V-ATPase subunits in *Diaphorina citri* (Hemiptera: Liviidae; NCBI:txid121845) as a step towards the development of future management strategies to reduce the psyllid vector of *Candidatus Liberibacter asiaticus* (CLas), the causative bacterial agent of Huanglongbing (HLB), also known as citrus greening disease.

METHODS

V-ATP synthase insect orthologs from *Acyrtosiphon pisum* (pea aphid) were obtained from the Kyoto Encyclopedia of Genes and Genomes (KEGG) database (RRID:SCR_012773). Additional ortholog subunits occurring in non-insect eukaryotes, like human (*Homo sapiens*), were obtained from the HUGO Gene Nomenclature Committee (HGNC) (RRID:SCR_002827) and the nonredundant National Center for Biotechnology Information (NCBI) Reference Sequence (Ref-Seq) database [8]. V-ATPase protein sequences were used to query the predicted protein set from the *D. citri* MCOT (Maker (RRID:SCR_005309), Cufflinks (RRID:SCR_014597), Oases (RRID:SCR_011896), and Trinity (RRID:SCR_013048)) transcriptome via protein BLAST (BLASTp) [9]. Reciprocal BLASTp analysis was performed to validate the *D. citri* MCOT significant hits using the NCBI nonredundant protein database [8]. *D. citri* V-ATPase genes were identified in the genome (version 1.91) by searching for the identified mapped MCOT models in the WebApollo (RRID:SCR_005321) system hosted at the Boyce Thompson Institute. Multiple alignments of the predicted *D. citri* MCOT proteins, other gene model sequences, and insect V-ATPase orthologs were performed using the European Bioinformatics Institute MUSCLE alignment online tool (RRID:SCR_004727) [10]. Further analysis using RNA sequencing (RNA-seq) reads, Illumina DNA sequencing reads, StringTie models, and Pacific Biosciences Isoform sequencing transcripts were used to manually annotate the final V-ATPase gene models. Manually annotated V-ATPase gene models were then integrated into the version 3.0 official gene set (OGS). V-ATPase genes were verified in WebApollo through analysis using *de novo*-assembled transcripts, Iso-seq transcripts, Augustus models, Mikado transcriptome, SwissProt proteins, and SNAP prediction models. A list of annotated *D. citri* identifiers and a sample of evidence supporting the annotated models is found in Table 1. A more detailed description of the annotation workflow is available (Figure 1) [11]. V-ATPase nomenclature is somewhat inconsistent in the literature and between species, therefore, we have used nomenclature standards reported in previous work on other Hemiptera [12, 13].

Reciprocal BLASTp of manually annotated v3.0 V-ATPase genes were performed at the NCBI comparing the Insecta taxid. Insect orthologs from *Acyrtosiphon pisum* (pea aphid) [14, 15], *Bemisia tabaci* (whitefly) [16], *Aedes aegypti* (yellow fever mosquito) [17], *Apis mellifera* (honeybee) [18], *Tribolium castaneum* (red flour beetle) [19], and *Drosophila*

Dec 17, 2020

Run

Bookmark

Copy / Fork

Annotating genes in *Diaphorina citri* genome version 3

Teresa D Shippy¹, S Miller², C Massimino³, C Vosburg [Indian River State College⁴, PS Hosmani⁵, M Flores-Gonzalez⁵, LA Mueller⁵, WB Hunter⁶, JB Benoit⁷, SJ Brown¹, T D'elia³, Surya Saha⁵

¹Kansas State University; ²Kansas State University, Allen County Community College; ³Indian River State College; ⁴Indian River State College, The Pennsylvania State University; ⁵Boyce Thompson Institute; ⁶USDA-ARS U.S. Horticultural Research Laboratory; ⁷University of Cincinnati

2 Works for me dx.doi.org/10.17504/protocols.io.bniimcce

D. citri annotation

Teresa Shippy

Kansas State University

Figure 1. Protocols.io protocol for psyllid genome curation [11]. <https://www.protocols.io/widgets/doi?uri=dx.doi.org/10.17504/protocols.io.bniimcce>

Table 1. Evidence for gene annotations. MCOT transcriptome identifiers included, if applicable.

Gene	Identifier	MCOT	de novo transcriptome	Iso-seq	RNA-seq	Ortholog
V-ATPase a1	Dcitr07g04330.1.1	MCOT05340.0.CO	X	X	X	X
V-ATPase a2	Dcitr07g01670.1.1	MCOT20572.0.CT	X	X	X	X
V-ATPase b	Dcitr12g08560.1.1	-	X	X	X	X
V-ATPase c	Dcitr06g11110.1.1	-	X	X	X	X
V-ATPase d	Dcitr04g06930.1.1	-	-	X	X	X
V-ATPase e	Dcitr03g19730.1.1	-	X	X	X	X
V-ATPase A	Dcitr06g09110.1.1	MCOT04747.0.CC	-	X	X	X
V-ATPase B	Dcitr09g08730.1.1	-	X	X	X	X
V-ATPase C	Dcitr02g01535.1.1	-	X	X	X	X
V-ATPase D	Dcitr09g02030.1.1	-	-	X	X	X
V-ATPase E	Dcitr04g09575.1.1	-	-	X	X	X
V-ATPase F	Dcitr07g06920.1.1	MCOT14638.0.CC	X	X	X	X
V-ATPase G	Dcitr11g08810.1.1	MCOT22289.0.CT	-	X	X	X
V-ATPase H (partial, N-terminus)	Dcitr00g06320.1.1	-	-	-	-	-
V-ATPase H (partial, C-terminus)	Dcitr01g01240.1.1	MCOT00604.0.CT	-	X	X	X
V-ATPase Ac45	Dcitr09g09620.1.1	MCOT16252.0.CC	X	X	X	X

MCOT: Maker, Cufflinks, Oases, Trinity.

melanogaster (fruit fly) [20] were obtained by reciprocal BLASTp (RRID:SCR_004870) analysis of the nonredundant protein database at NCBI [8]. A neighbor-joining phylogenetic tree was constructed using MUSCLE (RRID:SCR_011812) multiple sequence alignment with the Poisson correction method and 1000 replicate bootstrap test using full-length protein sequences in MEGA version 7 (RRID:SCR_000667) for the transmembrane complex, the catalytic complex, and the accessory subunit Ac45, respectively (Figures 2–4) [21]. The sequence accession numbers used in these analyses can be found in Tables 2–4. Comparative expression levels of *D. citri* V-ATPases throughout egg, nymph, and adult life stages in *D. citri* insects both exposed and not exposed to CLAs were determined using RNA-seq data and the Citrus Greening Expression Network (CGEN) [9]. These gene expression levels were visualized using the pheatmap package in R (RRID:SCR_016418) [22, 23]. Expression values for all samples discussed in this manuscript are visualized in Figures 5 and 6 and are reported as transcripts per million (TPM) in Table 5.

Figure 2. Phylogenetic analysis of V-ATPase V_0 transmembrane domain, subunits a–e. The tree was constructed with MEGA7 software [21] using MUSCLE for alignment of amino acid sequences, followed by neighbor-joining analysis with 1000 bootstrap replications. Values greater than 50 are shown at nodes. *Diaphorina citri* is marked in bold and color-coding indicates specific V_0 subunit groups. NCBI accession numbers are shown in Table 2.

Figure 3. Phylogenetic analysis of V-ATPase V₁, catalytic domain, subunits A-G. The tree was constructed with MEGA7 [21] software using MUSCLE for alignment of amino acid sequences, followed by neighbor-joining analysis with 1000 bootstrap replications. Values greater than 50 are shown at nodes. *Diaphorina citri* is marked in bold and color-coding indicates specific V₁ subunit groups. *D. citri* V-ATPase H (V₁-H) was annotated as two partial gene models, therefore, only the C-terminus, partial amino acid sequence of V₁-H was included in this analysis. NCBI accession numbers are shown in Table 3.

Figure 4. Phylogenetic analysis of V-ATPase accessory subunit Ac45. The tree was constructed with MEGA7 software [21] using MUSCLE for alignment of amino acid sequences, followed by neighbor-joining analysis with 1000 bootstrap replications. Values greater than 50 are shown at nodes. *Diaphorina citri* is marked in bold. NCBI accession numbers are shown in Table 4.

DATA VALIDATION AND QUALITY CONTROL

Genes encoding all 13 subunits required to build a single V-ATP synthase enzyme, as well as an accessory subunit S1 (*Ac45*) gene, were annotated in *D. citri*. There were no additional subunits found in *D. citri*, as reported in other metazoans [2]. Although insect V-ATPases are known to contain 13 subunits, there is variation in the gene copy number for individual subunits among different species (Tables 6–8). The V_0 transmembrane domain subunits *V-ATPase a*, *b*, and *e*; the V_1 catalytic domain subunits *V-ATPase A*, *C*, *D*, and *G*; and *Ac45*, all show variation in copy number among different species. The three Hemipterans analyzed (*D. citri*, *A. pisum*, and *B. tabaci*) maintain the same paralog number for all *V-ATPase* genes, except for *A. pisum V-ATPase D* and *G*, compared with the other orders (Table 7). This variation in copy number is interesting in contrast to the genes *V-ATPase c*, *d*, *B*, *E*, *F*, and *H*, which maintain only one gene copy across all orders of Insecta found in Tables 6–8. V-ATPase subunits have been studied in plants, animals, fungi, and insects, and certain genes have been highlighted for their functional versatility in serving cell needs. For example, yeast and mammals have numerous copies and isoforms of the transmembrane proteolipid *V-ATPase a* with functions supporting vacuoles, Golgi, neurons, osteoclasts, and epididymal cells [2]. *D. citri*, along with *A. pisum* and *B. tabaci*, has two copies of the *V-ATPase a* gene, whereas *D. melanogaster* has five copies (Table 6). In *D. citri*, a paralog of *V-ATPase a* was found, and they maintain differences in their amino acid sequences (Tables 2, 6) [24]. Phylogenetic analysis of V_0 subunit protein sequences supports that the

Figure 5. Comparative expression levels of the *Diaphorina citri* V-ATPase genes encoding the V_0 , transmembrane, subunits in *D. citri* insects reared on various infected and uninfected citrus varieties. Expression data were collected from the Citrus Greening Expression Network [9], with psyllid RNA-seq data from NCBI BioProjects PRJNA609978 and PRJNA448935, in addition to several published datasets [25–29]. Citrus hosts are abbreviated as *Csin* (*Citrus sinensis*), *Cmed* (*Citrus medica*), *Cret* (*Citrus reticulata*), and *Cmac* (*Citrus macrophylla*). Transcripts per million (TPM) values are listed in Table 5. Rows of genes and columns of RNA-seq data are clustered based on expression differences. (a) Expression data scaled by sample. (b) Expression data scaled by gene.

duplication event occurred before the divergence of Hemimetabola and Holometabola (Figure 2).

We identified complete genes in genome v3.0 for all the subunits except *V-ATPase H*. Using genome-independent transcript sequences [24], we were able to determine that the 3' portion of the *V-ATPase H* gene is located on chromosome 1, but the 5' end of the gene is on

Figure 6. Comparative expression levels of the *Diaphorina citri* V-ATPase genes encoding the V₁, catalytic, and accessory subunits in *D. citri* insects reared on various infected and uninfected citrus varieties. V-ATPase H was annotated as two partial models and are both represented here separately as V-ATPase H (5'), denoting the 5-prime end of the gene, and V-ATPase H (3'), denoting the 3-prime end of the gene. Expression data were collected from the Citrus Greening Expression Network [9], with psyllid RNA-seq data from NCBI BioProjects [PRJNA609978](#) and [PRJNA448935](#), in addition to several published datasets [25–29]. Citrus hosts are abbreviated as *Csin* (*Citrus sinensis*), *Cmed* (*Citrus medica*), *Cret* (*Citrus reticulata*), and *Cmac* (*Citrus macrophylla*). Transcript per million (TPM) values are listed in Table 5. Rows of genes and columns of RNA-seq data are clustered based on expression differences. (a) Expression data scaled by sample. (b) Expression data scaled by gene.

Table 2. *V-ATPase* transmembrane subunit (V_0) BLAST table.

Gene		<i>A. pisum</i>	<i>B. tabaci</i>	<i>T. castaneum</i>	<i>D. melanogaster</i>	<i>A. aegypti</i>	<i>A. mellifera</i>
<i>V-ATPase a1</i>	Accession	XP_029343960.1	XP_018903507.1	XP_008200952.1	NP_650722.1	XP_021706364.1	XP_006565533.1
	Bit score	1332	1312	1128	1151	1217	1166
	QC (%)	99	100	100	99	99	99
	Identity (%)	75.21	72.63	66.59	66.23	69.75	68.56
<i>V-ATPase a2</i>	Accession	XP_008183003.1	XP_018913655.1	XP_008200806.1	NP_733274.1	XP_021693139.1	XP_026298707.1
	Bit score	1369	1334	1331	1286	1369	1392
	QC (%)	100	100	100	100	100	100
	Identity (%)	79.10	76.42	76.18	76.19	77.21	79.05
<i>V-ATPase b</i>	Accession	NP_001155679.1	XP_018909463.1	NP_001161226.1	NP_001247111.1	XP_001662256.1	XP_392599.1
	Bit score	294	283	231	270	275	300
	QC (%)	99	99	98	99	99	99
	Identity (%)	73.79	73.30	63.05	72.95	74.4	76.33
<i>V-ATPase c</i>	Accession	NP_001155531.1	XP_018897791.1	XP_967959.1	NP_476801.1	XP_001654757.1	NP_001011570.1
	Bit score	268	250	258	271	267	259
	QC (%)	98	99	98	99	98	98
	Identity (%)	92.11	86.93	89.54	92.16	92.76	89.47
<i>V-ATPase d</i>	Accession	NP_001191854.1	XP_018903442.1	XP_974905.1	NP_570080.1	XP_001661299.1	XP_393438.2
	Bit score	656	669	671	661	677	679
	QC (%)	99	100	10	99	100	100
	Identity (%)	89.60	91.95	91.67%	90.20	91.95	92.82
<i>V-ATPase e</i>	Accession	XP_003242132.1	XP_018909271.1	XP_971898.1	NP_001097499.1	ABF18129.1	XP_624787.1
	Bit score	127	119	123	106	133	127
	QC (%)	96	96	94	96	100	94
	Identity (%)	69.51	70.73	72.50	58.54	70.59	71.25

Accession number, bit score, query coverage (QC), and identity results from protein BLAST analysis of annotated *D. citri* *V-ATPase* transmembrane subunit genes to their putative orthologs.

one of the unplaced chromosomes that make up chromosome 0 (Table 1). Tables 2–4 show the results of protein BLAST analysis comparing the same insects as found in Tables 6–8. Other than *Ac45*, all subunits share a relatively high sequence identity, approximately 57–94%, among individual pairwise alignments with each *D. citri* sequence (Tables 2–3). BLAST results of annotated gene models had high query coverage to orthologs, supporting the completeness of the annotated gene models. In contrast, the sequence identities of *Ac45*, approximately 24–33%, show the highest divergence when comparing *D. citri* to other insects (Table 4). For the V_0 , transmembrane domain, subunits in Table 2, proteolipid subunit c (*V-ATPase c*) maintains some of the highest percentages of sequence identity, highlighting the importance of the protein function to form the c-ring that rotates and ultimately translocates protons across various membranes [2]. This is supported by the data shown in Table 6, in which a single gene copy for *V-ATPase c* is maintained across different orders of insects.

The Citrus Greening Expression Network (CGEN) was used to compare transcript expression levels in various regions of *D. citri* that have either been exposed to or not exposed to CLas infection [9, 30, 31]. Figure 5 shows a heatmap comparing V_0 subunit expression levels found under various conditions. *V-ATPase c* is visually differentiated from other *V-ATPase* transmembrane subunit genes by its inflated expression levels (Figure 5a). *V-ATPase c* expression also shows a 2.63-fold increase, from 647.9 to 1705.25 TPM, in the guts of adult psyllids fed on infected *versus* uninfected *Citrus medica* leaves (Figure 5b, Table 5). These expression levels, coupled with the fundamental cellular nature and relatively even occurrence of *V-ATPases*, suggest that *V-ATPase* genes are good candidates for RNAi.

Table 3. V-ATPase catalytic subunit (V₁) BLAST table.

Gene		<i>A. pisum</i>	<i>B. tabaci</i>	<i>T. castaneum</i>	<i>D. melanogaster</i>	<i>A. aegypti</i>	<i>A. mellifera</i>
V-ATPase A	Accession	XP_008179407.1	XP_018897790.1	XP_976188.1	NP_001246015.1	XP_021709029.1	XP_016769524.1
	Bit score	1176	1187	1167	1162	1168	1174
	QC (%)	99	99	99	99	100	99
	Identity (%)	91.84	92.47	90.21	90.18	90.57	90.72
V-ATPase B	Accession	XP_003246082.1	XP_018896879.1	XP_967844.1	NP_001163597.1	XP_001651458.1	XP_624112.1
	Bit score	966	976	966	966	968	955
	QC (%)	98	99	98	98	98	98
	Identity (%)	93.88	94.33	93.48	93.88	94.09	92.87
V-ATPase C	Accession	XP_001946227.1	XP_018915661.1	XP_008195426.1	NP_477266.1	XP_021695404.1	XP_006562159.1
	Bit score	702	653	671	645	671	648
	QC (%)	99	99	98	98	99	9
	Identity (%)	87.05	82.81	82.20	79.11	81.77	79.95%
V-ATPase D	Accession	NP_001119691.1	XP_018904914.1	XP_975872.1	NP_651987.1	XP_001660426.1	XP_394769.2
	Bit score	420	423	426	408	401	404
	QC (%)	100	100	100	100	100	100
	Identity (%)	87.24	86.42	86.53	82.52	80.89	86.94
V-ATPase E	Accession	NP_001155650.1	XP_018901912.1	XP_970621.1	NP_001287182.1	XP_001655825.1	XP_625098.1
	Bit score	301	336	341	325	335	336
	QC (%)	99	100	100	100	100	100
	Identity (%)	73.21	71.68	73.01	71.24	70.80	73.01
V-ATPase F	Accession	NP_001119690.1	XP_018905603.1	XP_975016.1	NP_476969.1	XP_001655376.1	XP_624852.1
	Bit score	223	227	216	217	216	230
	QC (%)	98	98	98	98	100	98
	Identity (%)	85.25	88.52	82.79	81.15	82.26	88.52
V-ATPase G	Accession	NP_001119628.1	XP_018908074.1	XP_973974.1	NP_001287407.1	XP_001652605.1	XP_624346.1
	Bit score	184	188	173	154	171	180
	QC (%)	99	100	98	96	96	98
	Identity (%)	80.34	81.51	74.14	65.79	73.68	76.72
V-ATPase H (partial, N-terminus)	Accession	XP_001949116.3	XP_018904009.1	NP_001280516.1	NP_001260510.1	XP_001652018.1	XP_003251675.1
	Bit score	132	162	132	110	130	142
	QC (%)	90	99	99	84	91	94
	Identity (%)	57.69	68.42	57.26	56.70	58.10	64.22
V-ATPase H (partial, C-terminus)	Accession	XP_001949116.3	XP_018904009.1	NP_001280516.1	NP_001260510.1	XP_001652018.1	XP_003251675.1
	Bit score	530	506	513	493	498	490
	QC (%)	100	99	100	100	100	99
	Identity (%)	83.95	82.33	83.28	77.59	78.93	82.15
Accession number, bit score, query coverage (QC), and identity results from protein BLAST analysis of annotated <i>Diaphorina citri</i> V-ATPase catalytic subunit genes to their putative orthologs.							

Table 4. V-ATPase Ac45, accessory subunit, BLAST table.

Gene		<i>A. pisum</i>	<i>B. tabaci</i>	<i>T. castaneum</i>	<i>D. melanogaster</i>	<i>A. aegypti</i>	<i>A. mellifera</i>
Ac45	Accession	NP_001162140.1	XP_018899028.1	XP_974187.2	NP_610470.1	XP_001658652.1	XP_001121483.3
	Bit score	136	177	114	77.4	92.4	88.6
	QC (%)	83	97	100	96	95	93
	Identity (%)	30.87	33.49	27.03	24.94	23.81	26.35
Accession number, bit score, query coverage (QC), and identity results from protein BLAST analysis of annotated <i>Diaphorina citri</i> V-ATPase Ac45 gene to its putative orthologs.							

Silencing a V₀ transcript should have inhibitory effects on the assembly of the V-ATPase enzyme. In particular, if infected psyllids increase in their demand for higher V-ATPase *c* expression levels overall, knocking this transcript down is likely to be detrimental for the

Table 5. Expression values listed as transcripts per million (TPM), visualized in Figures 5 and 6.

Gene name	V-ATPase a1	V-ATPase a2	V-ATPase b	V-ATPase c	V-ATPase e	V-ATPase A	V-ATPase B	V-ATPase C	V-ATPase D	V-ATPase F	V-ATPase G	V-ATPase H (5')	V-ATPase H (3')	V-ATPase Ac45
Gene ID	Dcitr07 g04330.1.1	Dcitr07 g01670.1.1	Dcitr12 g08560.1.1	Dcitr06 g11110.1.1	Dcitr03 g19730.1.1	Dcitr06 g09110.1.1	Dcitr09 g08730.1.1	Dcitr02 g01530.1.1	Dcitr09 g02030.1.1	Dcitr07 g06920.1.1	Dcitr11 g08810.1.1	Dcitr00 g06320.1.1	Dcitr01 g01230.1.1	Dcitr09 g09620.1.1
Egg <i>Cmac</i> CLas- whole body	27.02	128.27	134.58	1085.07	312.06	343.23	15.59	49.46	567.45	199.16	698.5	161.78	248.69	148.17
Nymph <i>Cmed</i> CLas+ Low inf whole body	12.02	96.32	40.57	398.66	133.64	126.32	0	43.02	1.12	124.84	416.8	119.07	160.75	0
Nymph <i>Csin</i> CLas+ High inf whole body	8.82	89.21	39.09	334.65	131.69	102.13	0	36.49	1.4	117.74	408.72	99.61	144.58	0
Nymph <i>Csin</i> CLas- whole body	12.99	99.94	49.5	312.43	161.29	134.19	0	51.94	2.57	176.32	539.91	104.1	155.61	0
Nymph <i>Cmac</i> CLas- whole body	21.77	98.95	82.02	340.27	83.11	189.27	11.78	37.44	164.67	97.37	293.69	91	191.48	69.04
Nymph Citrus CLas- whole body	20.57	58.34	42.83	399.58	83.86	168.7	28.72	58.81	80.9	223.67	439.82	39.64	73.2	81.35
Nymph Citrus CLas+ whole body	4.94	38.29	59.33	621.98	146.78	160.92	2.44	53.38	190.77	444.42	766.12	44.97	115.28	114.6
Adult <i>Cmed</i> CLas- gut	38.18	289.58	200.59	647.9	426.18	605.11	26.07	50.69	381.68	296.91	1044.09	341.38	473.57	332.39
Adult <i>Cmed</i> CLas+ gut	30.2	319.44	230.85	1705.25	428.81	659.96	27.78	60.44	529.68	352.93	1163.03	317.72	485.17	329.78
Adult <i>Cmed</i> CLas+ High inf whole body	30.47	131.75	59.8	406.4	151.3	171.64	0	55.4	3.76	118.88	477.59	125.92	210.48	0
Adult <i>Cmed</i> CLas+ Low inf whole body	20.42	129.81	62.36	433.76	165.49	170.73	0	43.57	3.03	137.14	519.39	105.76	181.58	0
Adult <i>Cmed</i> CLas- whole body	30.31	113.42	49.08	412.59	132.76	149.36	0	71.67	2.27	70.33	411.72	118.34	191.46	0
Adult <i>Cmac</i> CLas- whole body	26.08	162.08	102.83	676.39	273.28	230.43	31.71	38.6	306.37	142.53	456.71	128.13	200.35	122.02
Adult Citrus CLas- whole body	36.98	101.48	106.15	554.39	212.37	217.42	27.13	54.78	123.29	228.76	592.56	83.13	188.23	96.36
Adult Citrus CLas+ whole body	19.8	77.29	133.73	951.84	248.6	268.61	12.34	73.2	257.5	196.34	698.74	131.62	218.34	158.58
Adult Citrus CLas- midgut	103.17	399.83	371.99	2604.59	688.74	825.5	13.38	56.53	681.66	519.97	1032.33	309.77	460.52	594.4
Adult Citrus CLas+ midgut	55.24	188.63	261.09	1449.95	529.49	549.52	8.89	33.98	732.48	484.27	1170.81	195.92	304.17	521.08
Adult <i>Cret</i> CLas- Female abdomen	48.78	117.05	150.69	1014.5	302.79	479.37	2.55	23.5	322.17	327.16	646.63	220.46	324.63	224.43
Adult <i>Cret</i> CLas- Female antennae	9.84	59.74	202.33	1200.58	515.42	374.86	0.79	16.81	407.15	343.42	564.73	165.93	204.21	300.71
Adult <i>Cret</i> CLas- Female head	19.05	57.69	174.15	1299.64	416.69	357.16	1.65	23.05	452.84	326.87	606.78	173.26	258.3	321.53
Adult <i>Cret</i> CLas- Female leg	13.96	40.8	110.91	1106.21	357.39	299.85	0.39	7.36	339.68	315.91	501.89	214.39	217.88	269.99
Adult <i>Cret</i> CLas- Female terminal abdomen	10.27	28.28	80.92	857.87	355.37	274.98	0.46	15.78	334.95	305.78	513.38	72.89	219.72	273.18
Adult <i>Cret</i> CLas- Female thorax	45.76	77.36	118.42	971.53	318.3	310.44	1.83	18.82	327.81	228.02	352.69	154.87	226.17	201.78
Adult <i>Cret</i> CLas- Male abdomen	55.04	113.66	163.22	1121.03	329.87	476.84	2.13	25.6	308.86	333.54	580.91	251.62	353.14	203.71
Adult <i>Cret</i> CLas- Male antennae	21.34	88.88	202.61	1216.63	502.11	454.87	2.06	20.5	366.31	317.82	489.63	163.36	206.39	300.97
Adult <i>Cret</i> CLas- Male head	22.42	57.66	180.09	1215.49	397.1	382.37	1.84	22.9	427.39	340.13	591.69	156.61	223.47	333.35
Adult <i>Cret</i> CLas- Male leg	26.24	70.25	126.32	1144.43	386.63	368.02	2.26	24.38	312.31	317.42	503.46	216.72	245.72	225.41
Adult <i>Cret</i> CLas- Male terminal abdomen	23.66	34.11	94.32	636.58	176.52	226.85	0.57	19.13	234.07	190.22	375.51	109.22	137.98	154.37
Adult <i>Cret</i> CLas- Male thorax	38.48	81.1	106.97	826.51	307.15	309.28	1.82	17.83	289.69	205.61	340.62	133.14	229.48	217.37
Adult <i>Cret</i> CLas- Female antennae Wu et al. [26]	55.37	150.12	175.28	828.29	314.86	436.78	33.85	23.48	389.56	253.26	576.96	141.07	182.16	314.72

Table 5. (Continued)

Gene name	V-ATPase <i>a</i> 1	V-ATPase <i>a</i> 2	V-ATPase <i>b</i>	V-ATPase <i>c</i>	V-ATPase <i>e</i>	V-ATPase <i>A</i>	V-ATPase <i>B</i>	V-ATPase <i>C</i>	V-ATPase <i>D</i>	V-ATPase <i>F</i>	V-ATPase <i>G</i>	V-ATPase <i>H</i> (5')	V-ATPase <i>H</i> (3')	V-ATPase <i>Ac45</i>
Gene ID	Dcitr07 g04330.1.1	Dcitr07 g01670.1.1	Dcitr12 g08560.1.1	Dcitr06 g11110.1.1	Dcitr03 g19730.1.1	Dcitr06 g09110.1.1	Dcitr09 g08730.1.1	Dcitr02 g01530.1.1	Dcitr09 g02030.1.1	Dcitr07 g06920.1.1	Dcitr11 g08810.1.1	Dcitr00 g06320.1.1	Dcitr01 g01230.1.1	Dcitr09 g09620.1.1
Adult <i>Cret</i> CLas– Female terminal abdomen Wu et al. [26]	46.1	121.03	149.2	727.77	282.02	427.76	9.27	18.35	373.06	280.05	731.99	106.17	164.41	273.56
Adult <i>Cret</i> CLas– Male antennae Wu et al. [26]	48.09	190.97	202.67	923.05	349.45	524.4	8.74	35.65	448.15	268.8	602.45	147.99	218.22	369.43
Adult <i>Cret</i> CLas– Male terminal abdomen Wu et al. [26]	77.21	129.36	162.97	503.06	208.23	381.74	8.29	36.86	235.06	163.07	405.46	95.28	131.34	183.8

Comparative expression levels in transcripts per million (TPM) of the *Diaphorina citri* V-ATPase genes encoding the V-ATPase V_0 transmembrane, V_1 catalytic, and accessory subunits in *D. citri* insects reared on various infected and uninfected citrus varieties. V-ATPase *H* was annotated as two partial models and are both represented here separately as V-ATPase *H* (5'), denoting the 5-prime end of the gene, and V-ATPase *H* (3'), denoting the 3-prime end of the gene. Expression data were collected from the Citrus Greening Expression Network [9], with psyllid RNA-Seq data obtained from NCBI BioProjects PRJNA609978 and PRJNA448935, in addition to several published datasets [25–29]. Citrus hosts are abbreviated as *Csin* (*Citrus sinensis*), *Cmed* (*Citrus medica*), *Cret* (*Citrus reticulata*), and *Cmac* (*Citrus macrophylla*).

Table 6. Gene copy comparison of V_0 transmembrane subunit V-ATPase genes in *Diaphorina citri* and orthologous insect genes.

Insect	V-ATPase <i>a</i>	V-ATPase <i>b</i>	V-ATPase <i>c</i>	V-ATPase <i>d</i>	V-ATPase <i>e</i>
<i>Diaphorina citri</i> (Hemiptera)	2	1	1	1	1
<i>Acyrtosiphon pisum</i> (Hemiptera)	2	1	1	1	1
<i>Bemisia tabaci</i> (Hemiptera)	2	1	1	1	1
<i>Tribolium castaneum</i> (Coleoptera)	2	2	1	1	2
<i>Drosophila melanogaster</i> (Diptera)	5	2	1	1	4
<i>Aedes aegypti</i> (Diptera)	3	1	1	1	2
<i>Apis mellifera</i> (Hymenoptera)	2	1	1	1	2

Table 7. Gene copy comparison of V_1 catalytic subunit V-ATPase genes in *Diaphorina citri* and orthologous insect genes.

Insect	V-ATPase <i>A</i>	V-ATPase <i>B</i>	V-ATPase <i>C</i>	V-ATPase <i>D</i>	V-ATPase <i>E</i>	V-ATPase <i>F</i>	V-ATPase <i>G</i>	V-ATPase <i>H</i>
<i>Diaphorina citri</i> (Hemiptera)	1	1	1	1	1	1	1	1
<i>Acyrtosiphon pisum</i> (Hemiptera)	1	1	1	2	1	1	2	1
<i>Bemisia tabaci</i> (Hemiptera)	1	1	1	1	1	1	1	1
<i>Tribolium castaneum</i> (Coleoptera)	1	1	1	3	1	1	2	1
<i>Drosophila melanogaster</i> (Diptera)	2	1	1	3	1	1	1	1
<i>Aedes aegypti</i> (Diptera)	1	1	2	2	1	1	3	1
<i>Apis mellifera</i> (Hymenoptera)	1	1	1	2	1	1	1	1

Table 8. Gene copy comparison of the *Ac45* gene in *Diaphorina citri* and orthologous insect genes.

Insect	<i>Ac45</i>
<i>Diaphorina citri</i> (Hemiptera)	1
<i>Acyrtosiphon pisum</i> (Hemiptera)	1
<i>Bemisia tabaci</i> (Hemiptera)	1
<i>Tribolium castaneum</i> (Coleoptera)	2
<i>Drosophila melanogaster</i> (Diptera)	2
<i>Aedes aegypti</i> (Diptera)	2
<i>Apis mellifera</i> (Hymenoptera)	1

insect. However, it remains unknown at this time whether the elevated expression of V-ATPase *c* relative to other subunits in infected psyllids is associated with higher demand of these proteins in the cell. This should, therefore, be studied further in future research.

Of the V_1 , catalytic domain, subunit genes, *V-ATPase A* and *V-ATPase B* maintain the highest percentages of sequence identity, consistent with the importance of their function in containing ATP binding sites at the V-ATPase subunits A/B protein interface (Table 3) [32]. Apart from *D. melanogaster*, *V-ATPase A* and *B* also maintain single copies of these two genes across different orders of insects, supporting their conserved nature compared with other genes of this enzyme (Table 7). *V-ATPase A* shows much higher expression than *V-ATPase B* across each measured variable, and *V-ATPase G* shows the highest expression in this group overall (Figure 6a). Unlike *V-ATPase c*, no significant differential expression was observed between the guts of insects reared on infected *versus* uninfected citrus trees (Figure 6b). However, *V-ATPase B* does show a reverse correlation, with a decrease in expression from 28.72 to 2.44 TPM in the whole body of *D. citri* nymphs raised on uninfected *versus* infected *Citrus spp.* (Table 5). A similar expression pattern can be seen throughout many of the *V-ATPase* catalytic genes, which may infer an interaction between these genes and pathogen infection. This warrants further investigations (Table 5).

Figures 2, 3, and 4 depict phylogenetic analyses for the V_0 transmembrane and V_1 catalytic domains, and the Ac45 protein of V-ATPase, respectively. The individual V-ATPase subunits form clades, regardless of insect species. These clades also have the highest bootstrap values. This agrees with previous research that describes the enzyme as ancient and highly conserved. The evolution of V-ATPase has been analyzed for gene duplication and divergence from other ATP synthases, like F- and A-ATPase, which occur across the three domains of life [3]. Figures 2, 3, and 4 concur and suggest that the V-ATPase enzyme utilized in these insects existed in their common ancestor before they diverged into their respective species. The proteolipid subunit c and subunit d have the shortest branch lengths in Figure 2, consistent with the data shown in Tables 2 and 6, which depict this to be of the most conserved subunits. Subunit c, which is required to form the critical c-ring rotor of V-ATPase [2], and subunit d, which may play a role as part of the central rotor of the V-ATPase [32], appear to have diverged the least compared with the other transmembrane domain subunits and other insect species. In contrast, subunit e has diverged the most (Figure 3). This is consistent with the variable gene copy number observed across different orders of insects and the lower percentages of protein sequence identity seen in *D. citri* pairwise alignments (Tables 2, 6). In addition, the function of subunit e is still unknown for the transmembrane domain subunits [5].

Figure 4 shows the evolutionary relatedness of the *D. citri* Ac45 protein. It is a relatively new protein, critically associated with the assembly of a certain cell type V-ATPase, and is still being studied [6]. For this select group of insect species, Ac45 groups and forms a clade with the other hemipteran protein sequences (Figure 4). *Ac45* is a variable gene when comparing *V-ATPase* across the domains of life, a paralog variability that is also seen among different orders of insects (Table 8) [6, 33]. *Ac45* has diverged the most of all the V-ATPase subunits in *D. citri* compared with other insects. This divergence is seen in phylogenetic analysis, denoted with longer branch lengths (Figure 5), and is also supported in the values of the pairwise alignments, in which the protein shares very little sequence identity across the query lengths (Table 4). Perhaps it is experimentally beneficial that the Ac45 protein shows the least conservation with other insect orthologs. It may serve as a species-specific targeted approach to limiting the psyllid from vectoring the causative agent of citrus greening disease, while leaving related species unharmed and their ecology intact. However, Ac45 shows a markedly depressed transcription level compared with other

subunits (Figure 6a, Table 5). This likely reflects the limits in resolution with current whole RNA isolation and sequencing methods; nevertheless, it still indicates the relatively low total expression. The Ac45 protein has not been observed to exist in every cell type depending on the organism and so is not necessarily utilized by every V-ATPase in the psyllid [6]. Thus, the expression data agree with previously published research.

CONCLUSION

The V-ATPase is a fundamental enzyme that functions exclusively as an ATP-dependent proton pump in almost every eukaryotic cell. V-ATPase allows for the proper functioning of endosomes and the Golgi apparatus, and it generates a proton-motive force in organelles and across plasma membranes, which is utilized as a driving force for secondary transport processes [1]. Identification of these enzymes in the hemipteran, *D. citri*, provides a novel insect lineage for studies of insect evolution and biology, and may also provide potential targets for *D. citri*-specific molecular mechanisms for the management of HLB in citrus production systems [34–36]. *D. citri* shows no deviation in the expected copy numbers of each of the V-ATPase genes (Tables 6–8). The data collected from *D. citri* reveal consistency among the genes previously characterized as highly conserved, such as V-ATPase c, d, A, and B (Tables 2–4) [3, 32]. While expression data were not available for V-ATPase d, V-ATPase c shows comparatively high expression levels overall, and differential expression – 647.9 versus 1705.25 TPM – in the guts of adult psyllids fed on uninfected versus infected *C. medica* leaves (Figure 5, Table 5). Conversely, the Ac45 gene shows low expression throughout life stages and tissues compared with other V-ATPase genes; however, the highly divergent nature of this gene may serve as a species-specific targeted approach to psyllid control (Table 4, Figure 6).

In hemipterans, RNAi efficacy has been successfully demonstrated for psyllids, whitefly, and leafhoppers [34–41]; planthoppers [42, 43]; bedbugs [44]; and others [45–49]. RNAi specifically targeting the V-ATPases in hemipteran insects has been reported for the corn planthopper, *Peregrinus maidis* (Ashmead) (Hemiptera: Delphacidae) [12]; the corn leafhopper, *Dalbulus maidis* (Hemiptera: Cicadellidae) [13]; the brown planthopper, *Nilaparvata lugens* (Stål) (Hemiptera: Delphacidae) [42]; and the bedbug, *Cimex lectularius* L. (Hemiptera: Cimicidae) [44], resulting in increased mortality and reduced fecundity. Thus, the highly divergent nature of these gene sequences provides unique targets that may serve as species-specific targeting for RNAi approaches in the management of psyllid vectors and other hemipteran pests [50, 51].

REUSE POTENTIAL

The manually curated gene models generated through this *D. citri* community annotation project will be available as part of the official gene set version 3. Analysis of these data, including BLAST and expression profiling, can be conducted using the citrusgreening.org website [52] and Citrus Greening Expression Network (CGEN). The improved annotations presented in this study will facilitate experimental design to investigate the potential of V-ATPases as gene targets for therapies to control *D. citri*. Research considering differential expression patterns of V-ATPase transcripts in psyllids fed on CLas-infected plants should be conducted. Additional studies are also required to confirm the role of the Ac45 protein, as its divergent nature may provide novel and species-specific gene targets, potentially through the use of RNAi, to control psyllid populations and reduce the effects of pathogens such as CLas.

DATA AVAILABILITY

The datasets supporting this article are available in the *GigaScience* GigaDB repository [24].

EDITOR'S NOTE

This article is one of a series of Data Releases crediting the outputs of a student-focused and community-driven manual annotation project curating gene models and if required, correcting assembly anomalies, for the *Diaphorina citri* genome project [53].

DECLARATIONS

LIST OF ABBREVIATIONS

ADP: Adenosine diphosphate; ATP: Adenosine triphosphate; BLASTp: protein BLAST; CLas: *Candidatus Liberibacter asiaticus*; Cmac: *Citrus macrophylla*; Cmed: *Citrus medica*; Cret: *Citrus reticulata*; Csin: *Citrus sinensis*; HLB: Huanglongbing; MCOT: Maker, Cufflinks, Oasis, Trinity; NCBI: National Center for Biotechnology Information; QC: Query coverage; CGEN: Citrus Greening Expression Network; RNAi: RNA interference; TPM: transcripts per million; V-ATPase: Vacuolar (H⁺)-ATP synthase; V₀: V-ATPase noncatalytic transmembrane domain; V₁: V-ATPase catalytic cytoplasmic domain.

ETHICAL APPROVAL

Not applicable.

CONSENT FOR PUBLICATION

Not applicable.

COMPETING INTERESTS

The authors declare that they have no competing interests.

FUNDING

This work was supported by USDA-NIFA grants 2015-70016-23028, HSI 2020-38422-32252 and 2020-70029-33199. And NIH grant IdeA P20GM103418.

AUTHORS' CONTRIBUTIONS

WBH, SJB, TD, and LAM conceptualized the study; TD, SS, TDS, and SJB supervised the study; SJB, TD, SS, and LAM contributed to project administration; RG conducted investigation; PH, MF-G, and SS contributed to software development; PH, MF-G, SS, TDS, and JB developed methodology; SJB, TD, WBH, and LAM acquired funding; RG and CM prepared and wrote the original draft; TD, SJB, SS, TDS, WT, WBH and JB reviewed and edited the draft. All authors read and approved the final version of the manuscript.

ACKNOWLEDGMENT

We would like to thank Helen Wiersma-Koch (Indian River State College) for her assistance.

REFERENCES

- 1 Nelson N, Perzov N, Cohen A et al. The cellular biology of proton-motive force generation by V-ATPases. *J. Exp. Biol.*, 2000; 203: 89–95.
- 2 Maxson ME, Grinstein S, The vacuolar-type H⁺-ATPase at a glance – more than a proton pump. *J. Cell Sci.*, 2014; 127: 4987–4993.

- 3 Hilario E, Gogarten JP, The prokaryote-to-eukaryote transition reflected in the evolution of the V/F/A-ATPase catalytic and proteolipid subunits. *J. Mol. Evol.*, 1998; **46**: 703–715.
- 4 Badillo-Vargas IE, Rotenberg D, Schneweis BA et al. RNA interference tools for the western flower thrips, *Frankliniella occidentalis*. *J. Insect Physiol.*, 2015; **76**: 36–46.
- 5 Beyenbach KW, Wieczorek H, The V-type H⁺ ATPase: molecular structure and function, physiological roles and regulation. *J. Exp. Biol.*, 2006; **209**: 577–589.
- 6 Jansen EJ, van Bakel NH, Coenen AJ et al. An isoform of the vacuolar (H⁺)-ATPase accessory subunit Ac45. *Cell Mol. Life Sci.*, 2010; **67**: 629–640.
- 7 Dow JA, Davies SA, Guo Y et al. Molecular genetic analysis of V-ATPase function in *Drosophila melanogaster*. *J. Exp. Biol.*, 1997; **200**: 237–245.
- 8 National Center for Biotechnology Information (NCBI). Ref-Seq Non-redundant Protein Database. Bethesda, MD: NCBI. <https://www.ncbi.nlm.nih.gov/refseq/about/nonredundantproteins/>. Accessed 2019.
- 9 Flores-Gonzalez M, Hosmani PS, Fernandez-Pozo N et al. Citrusgreening.org: An open access and integrated systems biology portal for the Huanglongbing (HLB) disease complex. bioRxiv. 2019; <http://dx.doi.org/10.1101/868364>.
- 10 Madeira F, Park YM, Lee J et al. The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic Acids Res.*, 2019; **47**(W1): W636–W641.
- 11 Shippy T, Miller S, Massimino C et al. Annotating genes in *Diaphorina citri* genome version 3. protocols.io. 2020; <https://dx.doi.org/10.17504/protocols.io.bniimcce>.
- 12 Yao J, Rotenberg D, Afsharifar A et al. Development of RNAi methods for *Peregrinus maidis*, the corn planthopper. *PLoS One*, 2013; **8**(8): e70243.
- 13 Jones T-KL, Bernal JS, Medina RF, Assessing the Functionality of RNA Interference (RNAi) in the Phloem-feeding Maize pest *Dalbulus maidis*. bioRxiv. 2021; <http://dx.doi.org/10.1101/2021.09.29.462424>.
- 14 International Aphid Genomics Consortium. Genome sequence of the pea aphid *Acyrtosiphon pisum*. *PLoS Biol.*, 2010; **8**: e1000313.
- 15 Li Y, Park H, Smith TE et al. Gene Family Evolution in the Pea Aphid Based on Chromosome-Level Genome Assembly. *Mol. Biol. Evol.*, 2019; **36**: 2143–2156.
- 16 Chen W, Hasegawa DK, Kaur N et al. The draft genome of whitefly *Bemisia tabaci* MEAM1, a global crop pest, provides novel insights into virus transmission, host adaptation, and insecticide resistance. *BMC Biol.*, 2016; **14**: 110.
- 17 Matthews BJ, Dudchenko O, Kingan SB et al. Improved reference genome of *Aedes aegypti* informs arbovirus vector control. *Nature*, 2018; **563**: 501–507.
- 18 Elsik CG, Worley KC, Bennett AK et al. Finding the missing honey bee genes: lessons learned from a genome upgrade. *BMC Genom.*, 2014; **15**: 86.
- 19 Richards S, Gibbs RA, Weinstock GM et al. Tribolium Genome Sequencing Consortium. The genome of the model beetle and pest *Tribolium castaneum*. *Nature*, 2008; **452**: 949–955.
- 20 Thurmond J, Goodman JL, Strelets VB et al. FlyBase 2.0: the next generation. *Nucleic Acids Res.*, 2019; **47**: D759–D765.
- 21 Kumar S, Stecher G, Tamura K, MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.*, 2016; **33**: 1870–1874.
- 22 R Core Team. R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing, 2020.
- 23 Kolde R, Pheatmap: Pretty Heatmaps (Version 1.0.12). 2020; <https://cran.r-project.org/package=pheatmap>.
- 24 Grace R, Massimino C, Shippy TD et al. Supporting data for “Genomic identification, annotation, and comparative analysis of Vacuolar-type ATP synthase subunits in *Diaphorina citri*”. *GigaScience Database*. 2022; <http://dx.doi.org/10.5524/100994>.
- 25 Reese J, Christenson MK, Leng N et al. Characterization of the Asian Citrus Psyllid Transcriptome. *J. Genomics*, 2014; **2**: 54–58.
- 26 Wu Z, Zhang H, Bin S et al. Antennal and abdominal transcriptomes reveal chemosensory genes in the Asian citrus psyllid, *Diaphorina citri*. *PLoS One*, 2016; **11**: e0159372.

- 27 Kruse A, Fattah-Hosseini S, Saha S et al. Combining 'omics and microscopy to visualize interactions between the Asian citrus psyllid vector and the Huanglongbing pathogen *Candidatus Liberibacter asiaticus* in the insect gut. *PLoS One*, 2017; **12**: e0179531.
- 28 Vyas M, Fisher TW, He R et al. Asian citrus psyllid expression profiles suggest *Candidatus Liberibacter asiaticus*-mediated alteration of adult nutrition and metabolism, and of nymphal development and immunity. *PLoS One*, 2015; **10**: e0130328.
- 29 Yu H-Z, Li N-Y, Zeng X-D et al. Transcriptome analyses of *Diaphorina citri* midgut responses to *Candidatus Liberibacter asiaticus* infection. *Insects*, 2020; **11**: 171.
- 30 Hosmani PS, Flores-Gonzalez M, Shippy T et al. Chromosomal length reference assembly for *Diaphorina citri* using single-molecule sequencing and Hi-C proximity ligation with manually curated genes in developmental, structural and immune pathways. bioRxiv. 2019; <http://dx.doi.org/10.1101/869685>.
- 31 Duan Y, Zhou L, Hall DG et al. Complete genome sequence of citrus huanglongbing bacterium, "*Candidatus Liberibacter asiaticus*" obtained through metagenomics. *Mol. Plant Microbe Interact.*, 2009; **22**(8): 1011–1020.
- 32 Toei M, Saum R, Forgac M. Regulation and isoform function of the V-ATPases. *Biochemistry*, 2010; **49**: 4715–4723.
- 33 Forgac M. Vacuolar ATPases: rotary proton pumps in physiology and pathophysiology. *Nat. Rev. Mol. Cell Biol.*, 2007; **8**: 917–929.
- 34 Hunter WB, Clarke S-KV, Mojica AFS et al. Advances in RNA suppression of the Asian citrus psyllid vector and bacteria (huanglongbing pathosystem). In: Asian Citrus Psyllid: Biology, Ecology and Management of the Huanglongbing Vector. Wallingford: CABI, 2020; pp. 258–284.
- 35 Hunter WB, Cooper WR, Sandoval-Mojica AF et al. Improving Suppression of Hemipteran Vectors and Bacterial Pathogens of Citrus and Solanaceous Plants: Advances in Antisense Oligonucleotides (FANA). *Front. Agron.*, 2021; **3**: 48.
- 36 Hunter WB, Wintermantel WM. Optimizing Efficient RNAi-Mediated Control of Hemipteran Pests (Psyllids, Leafhoppers, Whitefly): Modified Pyrimidines in dsRNA Triggers. *Plants*, 2021; **10**(9): 1782.
- 37 Andrade EC, Hunter WB. RNAi feeding bioassay: development of a non-transgenic approach to control Asian citrus psyllid and other hemipterans. *Entomol. Exp. Appl.*, 2017; **162**: 389–396.
- 38 Li H, Guan R, Guo H et al. New insights into an RNAi approach for plant defence against piercing-sucking and stem-borer insect pests. *Plant Cell Environ.*, 2015; **38**(11): 2277–2285.
- 39 Liu X, Zou Z, Zhang C et al. Knockdown of the Trehalose-6-Phosphate Synthase Gene Using RNA Interference Inhibits Synthesis of Trehalose and Increases Lethality Rate in Asian Citrus Psyllid, *Diaphorina citri* (Hemiptera: Psyllidae). *Insects*, 2020; **11**(9): 605.
- 40 Luo Y, Chen Q, Luan J et al. Towards an understanding of the molecular basis of effective RNAi against a global insect pest, the whitefly *Bemisia tabaci*. *Insect Biochem. Mol. Biol.*, 2017; **88**: 21–29.
- 41 Yu X, Killiny N. RNA interference of two glutathione S-transferase genes, *Diaphorina citri* DcGSTe2 and DcGSTd1, increases the susceptibility of Asian citrus psyllid (Hemiptera: Liviidae) to the pesticides fenpropathrin and thiamethoxam. *Pest Manag. Sci.*, 2018; **74**(3): 638–647.
- 42 Li J, Chen Q, Lin Y et al. RNA interference in *Nilaparvata lugens* (Homoptera: Delphacidae) based on dsRNA ingestion. *Pest Manag. Sci.*, 2011; **67**(7): 852–859.
- 43 Li J, Wang X-P, Wang M-Q et al. Advances in the use of the RNA interference technique in Hemiptera. *Insect Sci.*, 2013; **20**(1): 31–39.
- 44 Basnet S, Kamble ST. RNAi-Mediated Knockdown of vATPase Subunits Affects Survival and Reproduction of Bed Bugs (Hemiptera: Cimicidae). *J. Med. Entomol.*, 2018; **55**(3): 540–546.
- 45 Cagliari D, Dias NP, Galdeano DM et al. Management of Pest Insects and Plant Diseases by Non-Transformative RNAi. *Front. Plant Sci.*, 2019; **10**: 1319.
- 46 Christiaens O, Smagghe G. The challenge of RNAi-mediated control of hemipterans. *Curr. Opin. Insect Sci.*, 2014; **6**: 15–21.
- 47 Eakteiman G, Moses-Koch R, Moshitzky P et al. Targeting detoxification genes by phloem-mediated RNAi: A new approach for controlling phloem-feeding insect pests. *Insect Biochem. Mol. Biol.*, 2018; **100**: 10–21.
- 48 Fletcher SJ, Reeves PT, Hoang BT et al. A Perspective on RNAi-Based Biopesticides. *Front. Plant Sci.*, 2020; **11**: 51.

- 49 Jain RG, Robinson KE, Fletcher SJ et al. RNAi-Based Functional Genomics in Hemiptera. *Insects*, 2020; **11**(9): 557.
- 50 Whyard S, Singh AD, Wong S. Ingested double-stranded RNAs can act as species-specific insecticides. *Insect Biochem. Mol. Biol.*, 2009; **39**(11): 824–832.
- 51 Christiaens O, Whyard S, Vélez AM et al. Double-Stranded RNA Technology to Control Insect Pests: Current Status and Challenges. *Front. Plant Sci.*, 2020; **11**: 451.
- 52 Citrus Greening Solutions. 2018; <https://citrusgreening.org/>.
- 53 Asian citrus psyllid community annotation series. GigaByte. 2022; https://doi.org/10.46471/GIGABYTE_SERIES_0001.